

Great Vineyards Program

Non-Bearing Vineyard Assessment Form 2


Second-leaf or Older
(VSP or Sprawl, Spur-Pruned)
Time period: Mid to Late Season

Vineyard Name: _____ Date: _____

Block Description: Cultivar _____ Rootstock _____ Age _____

Check Yes or No to condition that applies to >90% of item (unless other % indicated)

	Yes	No
Vineyard Capacity		
0 to 5% Missing or dead vines		
50-100% of fruiting wire filled with cordon (exclusive of missing vines)		
Vine Training		
Developing cordons formed from lateral shoots		
Developing cordons originate from buds within 4" below cordon wire		
Developing cordons are shoots between 5/16" and 1/2 inch diameter		
Developing cordons tied appropriately to the cordon wire		
Incomplete cordons have distal shoot trained to cordon wire		
Shoots trained as cordon extensions are not excessively shaded		
4-6 shoots per linear foot of canopy (on average)		
Fruit load (if any) is appropriate for the canopy size		
Vine Health		
Absence of Crown Gall		
Absence of powdery mildew, downy mildew, phomopsis		
Absence of Pierce's disease symptoms on leaves and shoots		
Absence of severe leaf damage from insects or deer		
Leaf color normal, no apparent drought stress injury		
Leaf color normal, no apparent nutrient deficiencies		
Vineyard Floor		
Weed growth short and covers ≤ 10% of in-row area		
Cover crop appropriate for soil and climate		
Cover crop provides minimal competition with vines		
Vineyard Business		
Performance-based (e.g., yield, brix, TA, etc.) contract for future crop		



<u>Score</u>	<u>Rating</u>
90-100%	Excellent
80-89%	Commendable